



TB-500

Weekly Dosing Guide

• Dosing Protocol •

Week 1-2 → **5 units** daily

Week 3-8 → **10 units** daily

Increase only if well tolerated.

• Progress Overview •

Weeks 1-2:

- Subtle skin tone improvement
- Reduced irritation

Weeks 3-6:

- Improved elasticity
- Enhanced healing response

Weeks 6-8:

- Visible regenerative support.

Results are gradual and cumulative.



Administration Guidelines

- Flexible timing
- Can be taken morning or evening
- ✓ **Set Realistic Expectations**

Consistency > Aggressive dosing.

Cycle Length:

- ✓ 6-8 weeks on
- ✓ 4 weeks off



BPC-157

Weekly Dosing Guide

♦ Dosing Protocol ♦

Weeks 1-2 → **5 units** Daily

Weeks 3-4 → **5-10 units** Daily

Weeks 5-8+ → **10 units**: 1-2x Daily

Increase only if well tolerated.

♦ What to Expect: ♦

Weeks 1-2:

- Reduced inflammation
- Pain and tightness begin to improve

Weeks 3-6:

- Improved mobility
- Tissue repair becomes noticeable

Weeks 6-8:

- Increased stability and function

Results are gradual and cumulative



Administration Guidelines

- Subcutaneous injection
- Rotate injection sites with each injection
- ✓ Use a new sterile syringe for every administration
- ✓ Clean vial top with alcohol prior to drawing

Cycle Length:

- ✓ 4-8 weeks on
- ✓ 2-4 weeks off

For Research & Educational Use Only
Not for human or veterinary use



Wolverine

Weekly Dosing Guide

• Dosing Protocol •

Weeks 1-2 → **10 units** Daily

Weeks 3-4 → **10-15 units** Daily

Weeks 5-8+ → **15-20 units** Daily

Increase only if well tolerated.

• What to Expect •

Weeks 1-2

- Reduced inflammation
- Pain and tightness begin to improve

Weeks 3-6

- Improved mobility
- Tissue repair becomes noticeable

Weeks 6-8

- Increased stability and function
- Results are gradual and cumulative.



Administration Guidelines

- Subcutaneous injection
- Rotate injection sites with each injection
- Use a new sterile syringe for every administration
- Clean vial top with alcohol prior to drawing

Cycle Length:

✓ 4-8 weeks on ✓ 2-4 weeks off

For Research & Educational Use Only

Not for human or veterinary use



KPV

Weekly Dosing Guide

Weeks 1-2 → **5-10 units** Daily

Weeks 3-4 → **10-20 units** Daily

Weeks 5-8+ → **20-30 units** Daily

What to Expect

Weeks 1-2

- Reduced inflammation
- Decreased irritation/flare ups

Weeks 3-6

- Improved gut or skin stability
- Reduced inflammatory response

Weeks 6-8

- Sustained immune balance
- Improved overall symptom control

Results are gradual and cumulative.



Administration Guidelines

- Subcutaneous injection
- Rotate injection sites with each injection
- Use a new sterile syringe for every administration
- Clean vial top with alcohol prior to drawing

Cycle Length:

- ✓ 4-8 weeks on
- ✓ 2-4 weeks off



NAD+ DOSING GUIDE

Weekly Dosing & Administration

Weeks 1-2 → **25-50 Units**, 2-3x per week per injection

Weeks 3-8 → **50-100 Units**, 2-3x per week per injection

Increase only if well tolerated.

WHAT TO EXPECT

• Weeks 1-2

- Improved mental clarity
- More stable daily energy
- Reduced afternoon fatigue

Some may notice mild fatigue or or flushing initially.

• Weeks 3-6

- Improved physical recovery
- Better endurance and stamina
- Enhanced cognitive focus

Energy becomes more consistent throughout the day.

• Weeks 6-8

- Improved overall recovery capacity
- Better stress tolerance
- Sustained cellular energy support



ADMINISTRATION GUIDELINES

- * Subcutaneous injection
- * Common areas: abdomen, flank, or thigh
- * Rotate injection sites with each injection
- * Use a **new** *isterile*. syringe for every administration
- * Clean vial top with alcohol prior to drawing.

For Research & Educational Use Only
Not for human or veterinary use